

Development of silver resistance: a focus on wound care

The extensive use of silver-based products has led to the presence of silver in a variety of environments, including healthcare scenarios. This has increased the risk of development of silver resistance in bacteria. The authors carried out a systematic scoping review to assess the current state of antimicrobial resistance to silver, summarise the studies that discuss the rapidly developing issue of silver resistance in microorganisms (including wound bacteria) and evaluate its implications for wound care. A total of 105 articles were included in the review, indicating that silver resistance is an increasing topic in wound care, particularly in wound-derived microorganisms, posing a significant and increasing issue. This review identified several alternate options to silver-based therapies, including those that use the physical properties of wound dressings rather than chemical antimicrobial agents. While silver is an important tool for treating wound infections, its use should be cautious and the development of new antimicrobial agents that rely on alternate modes of action should be developed. To ensure that there are effective antimicrobial therapies available for wound care in the future, more studies should be performed to improve understanding on silver resistance, and for the development of antimicrobial agents and treatments that minimise the development of resistance.

The extensive use of silver-based products has led the release and accumulation of silver in river, soil and other environments (Kale et al, 2021), this makes it much more likely that bacteria will develop resistance to silver. The aim of this article is to undertake a scoping review to identify the present state of antimicrobial resistance (AMR) to silver, particularly in regard to wound care pathogens, and to discuss the implications of these findings in terms of current and future treatment options.

Silver in wound care-related healthcare

Infected wounds may require topical and/or systemic antimicrobial therapy (depending on the level of infection) (Wound Infection in Clinical Practice Working Group, 2008). However, the rise of multidrug-resistant bacteria has shifted the preference away from antibiotics towards topical antimicrobial agents, such as antiseptics to prevent and treat infections, including biofilms (Schultz et al, 2017).

The development and use of wound dressings that offer broad-spectrum antimicrobial activity has been significant with the use of several mechanisms of action being employed (Yousefian et al, 2023), including the use of silver (Khansa et al, 2019), iodine (Barreto et al, 2020), polyhexamethylene biguanide (PHMB) (Rippon et al, 2023), dialkylcarbamoyl chloride (DACC) (Totty et al, 2017; Rippon et

al, 2021) or medical grade (Manuka) honey (Scepankova et al, 2021). Antimicrobial wound dressings are widely used in wound care for their broad-spectrum antimicrobial properties (Yousefian et al, 2023). Among these, silver has been a primary choice for over two decades due to its effectiveness in combating infections and preventing biofilm formation (Frei et al, 2023).

Different forms of silver in wound care

Silver can be used in wound care in several different forms, including silver nitrate (a silver salt), silver sulfadiazine (a cream containing silver nitrate combined with sulfadiazine) and silver nanoparticles (AgNPs) (including nanocrystalline silver). Several silver-impregnated wound dressings have also been developed that provide silver to the wound. The role of AgNPs as an antimicrobial has gained significant attention due to their unique physiochemical properties that arise from the nanoscale dimensions of these particles (Rybka et al, 2022; Jangid et al, 2024). The positively charged silver ion (Ag^+) is the predominant form of silver that has antimicrobial activity (Lansdown, 2006; Percival et al, 2012), targeting microorganisms via several different modes of action (Summer et al, 2024). However, AgNPs can have direct antimicrobial activity by, for example, interacting with bacterial cell walls (Yin et al, 2020; Summer et al, 2024).

Mark G Rippon

Visiting Clinical Research Associate, University of Huddersfield, Huddersfield, UK

Alan A Rogers

Independent Wound Care Consultant, North Wales, UK

Key words

- Resistance
- Review
- Silver
- Wounds

Declarations

This study was funded by BSN medical GmbH (Essity Group), Schützenstraße 1-3, 22761 Hamburg, Germany.




Figure 1. Articles published per year and cumulatively related to silver resistance in bacteria between 1975 and 2023.

Development of silver resistance

Despite its widespread use, concerns have emerged regarding the long-term effects of environmental exposure and the potential development of bacterial resistance to silver (Ferdous and Nemmar, 2020). This development is worrisome, as indiscriminate use of silver compounds, similar to the overuse of antibiotics, could reduce their effectiveness in treating infections (McNeilly et al, 2021). AgNPs, while effective, accumulating evidence suggests that bacteria can show resistance to AgNPs (Kamat and Kumari, 2023; Li and Xu, 2024; Rodrigues et al, 2024).

Silver resistance associated with wound care

The overuse of silver (silver ions and nanoparticles) in commercially available healthcare products, including wound dressings, is a growing as a potential health concern due to the possible selection of tolerant or resistant bacteria, diverging from the once commonly held perception that bacteria could not develop resistance to silver (McNeilly et al, 2021), and silver resistance is an important issue for healthcare professionals (Hosny et al., 2019; McNeilly et al, 2021; Blackburn et al., 2023). Percival et al (2019) noted that there are only a limited number of studies documenting evidence of silver resistance in bacteria.

However, in a review of the emerging concern for silver resistance in microorganisms, McNeilly et al (2021) highlight that many important pathogenic bacteria show silver resistance (Gupta et al, 1999; Gunawan et al, 2013; Muller and Merrett, 2014; Panáček et al, 2018; Hosny et al, 2019; Valentin et al, 2020). Hosny et al (2019), in their study of silver resistance in clinical isolates from wounds and burns, note that their study "... is alarming regarding the spread of phenotypic silver-resistance ... in species where they were not

detectable before". Blackburn et al (2023) recently reviewed the evidence on AMR linked to antimicrobial use and observed that there is limited evidence to indicate that topical antimicrobials, including those containing silver, significantly contribute to resistance development. However, despite the limited clinical evidence, silver resistance is a critical issue for healthcare professionals, and the potential threat of resistance is a continuing essential area for exploration.

Silver resistance has become an increasingly critical area of research [Figure 1]. Recent research highlights the emergence of silver-resistant bacteria (Norton and Finley, 2021), the findings of which have been seen in studies in clinical settings where clinical isolates have been shown to exhibit silver resistance [Table 1].

Table 1. Studies demonstrating silver resistant in pathogens.

Microorganism	Form of silver	Reference
<i>P. aeruginosa</i> ¹	Silver nitrate	Muller and Merrett (2014)
	Silver nanoparticles	Panáček et al (2018); Liao et al (2019)
<i>S. aureus</i> ¹	Silver nitrate	Hosny et al (2019)
	Nanosilver	Valentin et al (2020)
<i>A. baumannii</i> ¹	Ionic silver	McNeilly et al (2021)
	Silver nanoparticles	McNeilly et al (2021)
<i>E. coli</i>	Ionic silver	Blanco Massani et al (2018)
	Silver nanoparticles	Panáček et al (2018)
<i>K. pneumoniae</i> ¹	Silver nitrate	Hanczvikkel et al (2018)
<i>Proteus</i> spp.	Nanosilver	Saeb et al (2017)

¹A WHO-designated critical priority pathogen that poses a high threat to public health characterised by resistance to multiple antibiotics (WHO, 2024)

This resistance could significantly impact current wound care practices, and the effectiveness of silver-containing medical devices, including silver wound dressings. For instance, laboratory analyses have demonstrated that some silver-impregnated dressings are less effective against resistant strains, suggesting the need for careful monitoring and judicious use of silver-based treatments (Sütterlin et al, 2012).

The issue of silver resistance aligns with broader concerns about AMR. Pathogens such as *E. coli*, *S. aureus*, *A. baumannii*, and *P. aeruginosa* have shown varying levels of silver resistance [Table 1]. These microorganisms are included in a group of important pathogens that are collectively referred to as the ESKAPE pathogens (Rice, 2008; Santajit and Indrawattana, 2016) and leading contributors to hospital-acquired infections and drug resistance (Ayobami et al, 2022; Bereanu et al, 2024).

A little-understood aspect of AMR is how heavy metals can impact antibiotic resistance. There are concerns of the coexistence of antibiotic and metal resistance: observational studies show that antibiotic-resistant bacteria are found at elevated levels in locations contaminated with metals (Gullberg et al, 2014). Antibiotic and metal resistance co-selection is where exposure to one selective agent allows adaptation to a second selective agent, i.e., where one antimicrobial (e.g., silver) selects for a resistance mechanism for both itself, and another antimicrobial (e.g., one or more antibiotics) (Gillieatt and Coleman, 2024). Three mechanisms have been suggested: co-resistance, where genes for resistance to different agents (e.g., silver and antibiotics) are on the same genetic element such as a plasmid, cross-resistance, where a single resistance mechanism (e.g., an efflux pump) provides resistance to multiple agents, and co-regulation, where the expression of multiple resistance genes is controlled by a shared regulatory system leading to simultaneous activation of resistance to different agents.

Implications of silver resistance in wound care

Given these developments, there is an urgent need to identify and utilize antimicrobial agents that are effective, non-toxic, and do not promote resistance. For example, DACC,

a relatively new and alternate antimicrobial treatment option in wound care, shows promise in meeting these criteria (Totty et al, 2017; Rippon et al, 2021; Rippon et al, 2023a). Unlike silver, which acts via chemical mechanisms, DACC works by physically disrupting microbial adhesion, reducing the risk of resistance, and promoting healing without cytotoxic or adverse effects. Such alternatives could play a crucial role in addressing the challenges posed by AMR.

Methods

A preliminary search of the literature in PubMed using the search strategy “silver” and “resistance” was conducted to assess the extent of the issue of silver resistance. A total of 9,531 references were identified from Jan-1975 (the first encountered silver-resistance was detected during an outbreak on a burns ward in 1975 (McHugh et al, 1975)) to end of 2023, with a further 852 articles being published between Jan-2024 and Oct-2024. Additionally, adding keywords “bacteria” and “microorganism” to the search strategy (to improve the relevance of the search) identified 5,502 references published in PubMed related to silver resistance in microorganisms. Note that, since 2020, there have been over 500 articles published per year [Figure 1] with, to Oct-2024, a further 514 articles have been published so far.

This scoping review was conducted in a semi-formal process and included several stages; (1) defining the research question; (2) identifying relevant studies; (3) study selection; (4) summarising and reporting the results. The research question was defined using the PEO framework [Table 2], which we considered an appropriate framework for our review (Doody and Bailey, 2016). The PEO framework breaks the topic of our review into three separate areas: the Population to focus on in the review, the Exposure (the issue of interest), and the Outcomes or themes to examine. Once the scope of the review was defined (to review the evidence for the development of silver resistance in microorganisms) relevant studies were identified. The PRISMA-ScR framework was used for reporting the results [Figure 2].

Research questions

Several research questions were identified as part of the development process for this review:

1. Is silver resistance a rapidly developing issue in microorganism populations?
2. Is silver resistance a rapidly developing issue in wound pathogens?
3. What are the implications of silver resistance for wound care?

Table 2. PEO framework to identify the research question.

Population	Patients and pre-clinical studies
Exposure	Wound infection
Outcome	Silver resistance

Search strategy and eligibility criteria

PubMed was searched using keywords “silver,” “resistant,” “resistance,” “bacteria,” “bacterial,” and “antimicrobial.” Articles published between January 1975 and October 15, 2024 were identified. Inclusion criteria for the search included primary evidence studies reporting silver resistance, laboratory, and clinical studies, with full-text articles written in the English language. Exclusion criteria included review articles, and papers not published in the English language.

Study selection, data extraction, and analysis

The process for study selection is outlined in the PRISMA–ScR flow diagram [Figure 2]. The titles and abstracts of each result was screened against the inclusion and exclusion criteria. Full texts of articles meeting the inclusion criteria were independently assessed by two authors (MR and AR). Following full-text screening, studies that met the inclusion criteria underwent data extraction. Collated information included the following data points: study aims and objectives; design and methodology; sample size; wound types; details of the outcome measures; and main study results [Table 3]. The primary outcome was the reporting of silver resistance in microorganisms.

Results

The findings were summarised and described narratively, under various collective headings based upon the research questions. The search identified 5,295 potential records, and records from other sources (e.g., hand searching) (n=9) were added, resulting in 5,304 records. Following review of titles and abstracts, 105 full text articles were retrieved and included in the narrative review [Figure 2]. An overview of the study characteristics is presented in Table 4.

Note that many articles were identified in the initial review that were related to the use of “silver” against “drug-resistant” microorganisms (which did not involve any assessment of silver resistance); this led to




Figure 2

them being captured by the search strategy. It was felt that adding any clarifying search keywords to try and reduce the number of these non-relevant articles may have impacted on capturing potentially relevant articles.

Discussion

Table 5 indicates the articles featuring ESKAPE pathogens (and other important microorganisms) reported to show phenotypic silver resistance.

Is silver resistance a rapidly developing issue in microorganism populations?

Considering the extensive use of silver ions as an antimicrobial in domestic, industrial, and medical applications (Ferdous and Nemmar, 2020), concerns have been raised over the potential for silver ion resistance emergence in bacteria of clinical relevance and to thereby compromise its therapeutic utility. Bacterial resistance to cationic silver (Ag+) has been

Figure 2. PRISMA–ScR flow diagram to indicate the search results based upon the search strategy.

Table 3. Main article assessment criteria.	
Criteria	Summary
Aims/objectives	Articles reporting on silver resistance
Design/methodology	Only reporting primary pre-clinical and clinical evidence
Wound type	Skin wounds (acute and chronic wounds)
Outcome measures	Outcome measures as set out in study methodology related to silver resistance in microorganisms
Main results	Results related to stated outcome measures

Table 4. Overview of characteristics of silver resistance studies.

Author	Study aim	Form of silver
Alhajjar et al (2022)	Adaptation of <i>E. coli</i> for resistance to Ag-NPs by repeated exposure to sub-MICs of the antimicrobials	Silver nanoparticles
Ali et al (2020)	Molecular basis of AgrR-mediated gene expression resulting in silver resistance	N/A
Alotaibi et al (2022)	Assessment of whether combinations of AgNPs and conventional antimicrobials produce synergistic responses	Silver nanoparticles
Andrade et al (2018)	Search for presence of acquired silver resistance genes in <i>E. cloacae</i> Complex and <i>E. aerogenes</i>	N/A
Annear et al (1976)	Demonstrate transferability of silver resistance in <i>E. cloacae</i> isolated from burn wound	Silver nitrate
Arrault et al (2023)	Assessment of solution structures of SilF in its free and Ag-bound forms	N/A
Arrieta-Gisasola et al (2025)	Detection of mobile genetic elements conferring heavy metal resistance in <i>Salmonella</i>	N/A
Ashraf et al (2014)	Efficacy of lysozyme-coated AgNPs for their antimicrobial activity against panel of bacteria	Silver nanoparticles
Asiani et al (2016)	Testing model of exporting Ag ⁺ in <i>Salmonella</i> via various sil operon components	N/A
Babel et al (2021)	Assessment of interplay between SilE and SilB peptides for silver resistance	N/A
Babu et al (2011)	Analysis of global transcriptome of <i>B. cereus</i> in response to silver nitrate stress	Silver nitrate
Bearson et al (2020)	Role of Salmonella Genetic Island 4 in metal tolerance of <i>S. enterica</i>	N/A
Billman-Jacobe et al (2018)	Assessment of pSTM6-275 plasmid of <i>S. enterica</i> to confer heavy metal resistance	N/A
Binsuwaidan et al (2024)	Effects of exposure to ionic silver in wound-associated bacterial pathogens	Silver nitrate
Brady et al (2003)	Assessment of silver disinfectant persistence to pathogenic bacteria on treated surfaces	Silver nitrate
Bridges et al (1979)	Emergence of silver-resistant <i>P. aeruginosa</i> in a burns unit	Silver nitrate
Chabert et al (2018)	Molecular analysis of structural folding of SilE model peptides	N/A
Chalmers et al (2018)	Assessment of genetic elements involved in heavy metal resistance in swine <i>E. coli</i>	N/A
Cidre et al (2017)	Assessment of genetic elements of resistance and their interplay	N/A
Clark et al (2020)	Distribution of heavy metal resistance elements in <i>Salmonella</i> populations	N/A
Dunne et al (2017)	Complete genomic sequence of <i>E. coli</i> strain detailing silver resistance genetic elements	N/A
Elarabi et al (2023)	Assessment of genome of <i>R. planticola</i> detailing heavy metal resistance elements	N/A
Elbehiry et al (2019)	Evaluation of resistance to AgNPs and AuNPs in <i>S. aureus</i>	Silver nanoparticles
Elkrewi et al (2017)	Assessment of prevalence of cryptic Ag ⁺ resistance amongst clinical isolates of Gram-negative bacteria	Silver nitrate
Finley et al (2015)	Identification of first strains of clinical bacteria expressing clinically relevant silver resistance	Silver nitrate
Foka et al (2020)	Genetic analysis and silver resistance gene detection in vancomycin-resistant <i>enterococci</i>	N/A

Table 4. Overview of characteristics of silver resistance studies.

Author	Study aim	Form of silver
Franke et al (2001)	Characterisation of silver resistance system in <i>E. coli</i>	N/A
Fuentes-Castillo et al (2021)	Genomic characterisation multidrug resistant <i>E. coli</i>	N/A
Gokulan et al (2017)	Assessment of specificity of silver resistance genes in <i>Salmonella Typhimurium</i>	Silver nitrate
González-Fernández et al (2020)	Assessment of antibacterial effect of silver nanorings compared with other AgNPs in silver-resistant <i>Salmonella</i>	Silver nanorings (AgNPs)
Graves et al (2015)	Characterisation of rapid evolution of AgNP resistance in <i>E. coli</i>	Silver nanoparticles
Grim et al (2013)	Genetic analysis of pathogen <i>Cronobacter spp.</i> including silver resistance genes	N/A
Gugala et al (2018)	Chemical genetic screen of <i>E. coli</i> to identify silver sensitive or resistant deletion strains	Silver nitrate
Gugala et al (2019)	Susceptibility of <i>E. coli</i> , <i>P. aeruginosa</i> and <i>S. aureus</i> to silver	Silver nitrate
Gullberg et al (2014)	Effects on low level antibiotics and heavy metals on selection of ESBL plasmid identified in <i>K. pneumoniae</i> and <i>E. coli</i>	Ionic silver
Guo et al (2019)	Assessment of the impact of AgNPs on <i>P. aeruginosa</i> biofilm	Silver nanoparticles
Gupta et al (1998)	Effect of halides on plasmid-mediated silver resistance in <i>E. coli</i>	Silver nitrate
Gupta et al (2001)	Assessment of diversity of silver resistance genes in plasmids	Silver nitrate
Gupta (1999)	Use RT-PCR to characterise to analyse transcripts that form long multi-gene operons	N/A
Haefeli et al (1984)	Characterisation of a silver-resistant <i>P. stutzeri</i> isolated from a silver mine	Silver nitrate
Håkonsholm et al (2023)	Characterisation of antibiotic- and heavy metal-resistance genes on plasmids	N/A
Håkonsholm et al (2022)	Whole-genome analysis to characterise resistance in <i>K. pneumoniae</i>	N/A
Hanczvikkel et al (2018)	Evolution of transmissible silver resistance in <i>K. pneumoniae</i>	Silver nitrate
He et al (2022)	Characterisation of hypervirulent carbapenem-resistant <i>K. pneumoniae</i>	N/A
Hosnedlova et al (2022)	Assessment of effect of AgNPs on <i>S. aureus</i> and <i>E. coli</i> biofilms	Silver nanoparticles
Hosny et al (2019)	Detection of silver resistance in isolates from wounds and burns and characterisation of plasmid-mediated silver resistance genes	Silver nitrate
Johnson et al (2005)	Genetic characterisation of a pathogenic <i>E. coli</i> plasmid	N/A
Joseph et al (2012)	Analysis of genomic sequences in <i>Cronobacter</i> species	N/A
Kamatthewatta et al (2020)	Detection of pathogen <i>E. hormaechei</i> and genomic analysis	N/A
Kaur and Vadehra (1986)	Characterisation of mechanism of silver resistance in <i>K. pneumoniae</i>	Silver nitrate
Kędziora et al (2020)	Assessment of long-term exposure of bacteria to silver nanoformulations	Silver nanoparticles
Khor and Jegathesan (1983)	Characterisation of heavy metal resistance in antibiotic-resistant Gram negative bacteria	Silver nitrate
Klonowska et al (2020)	Genomic analysis of heavy metal resistance genes in <i>C. neoaledonicus</i>	N/A

Table 4. Overview of characteristics of silver resistance studies.

Author	Study aim	Form of silver
Kremer and Hoffmann (2012)	Analysis of genetic differences in bacteria causing septicemia outbreak	Silver nitrate
Kucerova et al (2010)	Genomic analysis of <i>C. sakazakii</i> and silver resistance genes	N/A
Lau et al (2017)	Analysis of the effect of PVP-capped AgNPs on <i>Citrobacter sp.</i> and <i>Enterococcus sp.</i>	Silver nanoparticles
Li et al (1997)	Examination of the role of porin deficiency and its role in silver resistance	Silver nitrate
Lima de Silva et al (2012)	Assessment of heavy metal tolerance of sewage-isolated bacteria	Silver nitrate
Loh et al (2009)	Determination of the prevalence of silver resistance genes in bacteria isolates from humans and animals and their susceptibility to silver-containing Hydrofiber wound dressing	Silver dressing
Losasso et al (2014)	Assessment of antibacterial activity of AgNPs of different <i>Salmonella</i> serovars	Silver nanoparticles
Mann et al (2021)	Assessment of the long-term use of AgNPs on biofilm-forming pathogenic bacteria	Silver nanoparticles
Mastrorilli et al (2018)	Genomic analysis of <i>Salmonella</i> serovar including heavy metal resistance elements	N/A
McCarlie et al (2023)	Genomic analysis of highly resistant <i>Serratia sp.</i>	N/A
McNeilly et al (2023)	Assessment of AgNP adaptation in <i>A. baumannii</i>	Silver nanoparticles
Miloud et al (2021)	Assessment and genetic analysis of heavy metal and antibiotic resistance in bacteria	Silver nitrate
Monych and Turner (2020)	Characterisation of interaction between <i>P. aeruginosa</i> and tolerance of <i>S. aureus</i> to silver	Silver nitrate
Muller and Merrett (2014)	Analysis of pyocyanin production by <i>P. aeruginosa</i> and resistance to silver	Silver nitrate
Muller (2018)	Assessment of redox-based silver resistance gained by pyocyanin production by <i>P. aeruginosa</i>	Silver nitrate
Nicolás et al (2018)	Genetic analysis of clinical isolate of <i>K. quasipneumoniae subsp. similipneumoniae</i>	N/A
Panáček et al (2021)	Characterisation of resistance to AgNPs by binding AgNPs to cyanographene	Silver nanoparticles
Pant et al (2022)	Characterisation of overcoming of AgNP resistance in <i>P. aeruginosa</i> using bismuth nanoparticles	Silver nanoparticles
Percival et al (2008)	Prevalence of silver resistance in bacteria isolated from diabetic foot ulcers	Aquacel Ag
Pirnay et al (2003)	Molecular epidemiology analysis of silver-resistant <i>P. aeruginosa</i> clone	Silver sulfadiazine
Potgieter and Meidany (2018)	Evaluation of the penetration of nanocrystalline silver through wound dressing mediums	Nanocrystalline silver
Randall et al (2015)	Molecular and genetic analysis of silver resistance in <i>E. coli</i>	Silver nitrate
Riley and Mee (1982)	Assessment of the susceptibility of <i>Bacteoides spp.</i> to heavy metals	Silver nitrate
Saeb et al (2017)	Genetic analysis of a spontaneous nanosilver resistant <i>P. mirabilis</i> strain SCDRI	Silver nanoparticles
Safain et al (2023)	Assessment of prevalence of silver resistance determinants in wound infection bacteria	Silver nitrate

Table 4. Overview of characteristics of silver resistance studies.		
Author	Study aim	Form of silver
Salomoni et al (2017)	Antibacterial effect of AgNPs in <i>P. aeruginosa</i>	Silver nanoparticles
Sanjar et al (2024)	Genomic analysis of <i>P. aeruginosa</i> isolated from post-burn infections	N/A
Sano et al (2023)	Genomic analysis of <i>L. adecarboxylata</i>	N/A
Sedlak et al (2012)	Engineering of a silver-binding periplasmic protein promoting silver tolerance	Silver nitrate
Silver et al (1999)	Genetic analysis of silver resistance genes	Silver nitrate
Souza et al (2022)	Analysis of heavy metal and biocide resistance genes in <i>S. enterica</i>	N/A
Staehlin et al (2016)	Molecular and genetic analysis of heavy metal resistance in <i>Enterobacter</i>	N/A
Su et al (2011)	Assessment of novel nanohybrids of AgNPs on clay platelets for inhibiting silver resistance	Silver nanoparticles
Sütterlin et al (2017)	Genetic analysis of silver resistance genes in isolates of <i>Enterobacter</i> and <i>Klebsiella</i> species	N/A
Sütterlin et al (2014)	Genetic analysis of silver resistance genes in <i>E. coli</i> isolates	N/A
Sütterlin et al (2012)	Assessment of silver-based wound dressings on bacteria isolated from chronic leg ulcers and susceptibility in vitro to silver	Aquacel Ag; silver nitrate
Sütterlin et al (2018)	Genetic analysis of heavy metal susceptibility genes of <i>E. coli</i>	N/A
Vasishta et al (1989)	Assessment of heavy metal resistance to clinical isolates of <i>P. aeruginosa</i>	Silver nitrate
Vázquez et al (2023)	Genetic analysis of antibiotic resistance in clinical isolates of <i>S. enterica</i>	N/A
Vilela et al (2022)	Prevalence analysis of silver resistance elements in <i>S. enterica</i>	N/A
Villapún et al (2021)	Assessment of effects of repeated exposure of nosocomial pathogens to silver and selection of silver resistance	Silver nitrate
Wang et al (2022)	Genetic characterisation of silver and antibiotic resistance among Gram-negative pathogens isolated from wounds	N/A
Wiegand et al (2012)	Assessment of adaptation capacity of <i>S. aureus</i> to antiseptics (e.g., silver nitrate)	Silver nitrate
Woods et al (2009)	Genetic analysis of prevalence of silver resistance genes in bacteria isolated from wounds	Silver nitrate
Woolley et al (2022)	Molecular and genetic analysis of high level silver ion tolerance in <i>K. pneumoniae</i>	Silver nitrate
Woyda et al (2024)	Assessment of genetic characteristics of <i>Salmonella</i> isolates	N/A
Wu et al (2022)	Assessment of silver resistance development in bacteria challenged by AgNPs	Silver nanoparticles
Wu et al (2007)	Genomic analysis of effectiveness of silver to prevent biofilm formation	Silver nitrate
Yang et al (2020)	Assessment of antibiotic and heavy metal resistance in <i>E. coli</i>	Silver nitrate
Zhao and Kuipers (2021)	Assessment antimicrobial activity of Ag-nisin NPs in biofilm-forming bacteria	Silver nanoparticles; silver nitrate
Zingale et al (2023)	Molecular analysis of interaction of silver ions with SILE model peptides	Silver nitrate
Zingali et al (2020)	Genetic analysis of multiple drug resistant plasmid in <i>E. coli</i>	N/A

Table 5. Distribution of important silver-resistant microorganisms in articles featured in review.

	ESKAPE pathogens												
	<i>E. faecium</i>	<i>S. aureus</i>	<i>K. pneumoniae</i>	<i>A. baumannii</i>	<i>P. aeruginosa</i>	<i>Enterobacter spp.</i>	<i>B. subtilis</i>	<i>Salmonella spp.</i>	<i>E. coli</i>	<i>E. faecalis</i>	Others	<i>Candida spp.</i>	<i>S. epidermidis</i>
Alotaibi et al (2022)		■	■	■	■				■		■		■
Ashraf et al (2014)		■	■		■		■	■	■				
Binsuwaidan et al (2024)		■			■				■				
Brady et al (2003)	■	■	■			■		■	■	■			
Bridges et al (1979)					■								
Elbehiry et al (2019)		■											
Elkrewi et al (2017)			■	■	■	■			■		■		
Finley et al (2015)	■	■	■		■	■			■		■	■	
González-Fernández et al (2020)		■	■		■				■	■	■		■
Graves et al (2015)									■				
Guo et al (2019)					■								
Gugala et al (2019)		■			■				■				
Gupta et al (1998)									■				
Hanczvikkel et al (2018)			■										
Hosnedlova et al (2022)		■							■				
Hosny et al (2019)		■	■	■	■	■			■				
Kaur and Vadehra (1986)			■										
Kędziora et al (2020)		■	■			■			■				
Khor and Jegathesan (1983)			■		■	■		■	■		■		
Lau et al (2017)	■										■		
Lima de Silva et al (2012)			■	■	■	■					■		
Loh et al (2009)		■											
Losasso et al (2014)								■					
Mann et al (2021)					■								
McNeilly et al (2023)				■									
Miloud et al (2021)		■	■			■			■		■		

Table 5. Distribution of important silver-resistant microorganisms in articles featured in review.

	ESKAPE pathogens												
	<i>E. faecium</i>	<i>S. aureus</i>	<i>K. pneumoniae</i>	<i>A. baumannii</i>	<i>P. aeruginosa</i>	<i>Enterobacter</i> spp.	<i>B. subtilis</i>	<i>Salmonella</i> spp.	<i>E. coli</i>	<i>E. faecalis</i>	Others	<i>Candida</i> spp.	<i>S. epidermidis</i>
Monych and Turner (2020)		■											
Muller (2018)		■							■				
Muller and Merrett (2014)					■								
Panáček et al (2021)		■	■						■				
Pant et al (2022)					■								
Percival et al (2008)						■							
Pirnay et al (2003)					■								
Potgieter and Meidany (2018)		■			■						■		
Riley and Mee (1982)											■		
Salomoni et al (2017)					■								
Safain et al (2023)		■	■	■	■	■			■		■		
Sedlak et al (2012)									■				
Souza et al (2022)								■					
Su et al (2011)		■						■					
Sütterlin et al (2012)	■	■		■	■						■		
Sütterlin et al (2017)			■			■			■				
Sütterlin et al (2018)									■				
Vasishta et al (1989)					■								
Villapún et al (2021)		■			■				■				■
Wiegand et al (2012)		■											
Woods et al (2009)	■	■		■	■	■			■	■	■		
Wu et al (2007)									■				
Yang et al (2020)									■				
Zhao et al (2021)		■	■	■	■				■				

Notes:

■ – general reference to genus mentioned rather than specific species

Lighter grey references are ESKAPE pathogen-related articles; darker grey references are wound- and ESKAPE-related articles

recognized for several decades (Gupta et al., 1999), it was a common perception that resistance to nanosilver was unlikely, due to the multitargeting antimicrobial mechanisms of the nanoparticles (Gunawan et al, 2017; Valentin et al, 2020). There has been growing evidence regarding the development of adaptation phenomena to nanosilver, and this evidence has been observed in several bacterial species, including those of clinical significance, such as *E. coli*, *P. aeruginosa*, and *S. aureus* (Gunawan et al, 2013; Graves et al, 2015; Panáček et al, 2018; Valentin et al, 2020; Mann et al, 2021; Stabryla et al, 2021). The introduction of AgNP technologies and its effective use for treating wound infections have been suggested to reduce the likelihood of silver resistance (Pelgrift and Friedman, 2013). However, several articles have raised concerns over the development of silver resistance because of AgNP use (Kamat and Kumari, 2023; May et al, 2022).

The results from this review show that there is an increasing number of articles being published that are related to silver resistance in microorganisms, particularly in the last 5–10 years ($n=105$). Resistance to silver can be described as genotypic or phenotypic. Genotypic silver resistance refers to the inherent or acquired genetic mechanisms (determined by specific genes and plasmids) that allow bacteria to survive exposure to silver compounds (Randall et al, 2015). Phenotypic

silver resistance refers to a microorganism's observable ability to survive and grow in the presence of silver compounds, even though it doesn't involve any genetic changes (Corona and Martinez, 2013). Phenotypic resistance include specific processes such as biofilm formation. **Figure 3** shows the increase in articles in a subset of these 105 articles, namely, describing phenotypic silver resistance ($n=50$).

Wiegand et al (2012) analysed the adaptation capacity of *S. aureus* to several commonly used antiseptics (including silver nitrate). The bacteria were incubated with the appropriate IC50 concentrations of the antiseptics for 100 days. *S. aureus* quickly adapted to high concentrations of silver nitrate over time. The authors noted that, although silver-containing wound dressings were still as effective against silver-adapted *S. aureus*, overuse of silver may raise future problems. Elkrewi et al (2017) surveyed the prevalence of silver resistance in clinical isolates and reported that overt silver resistance is not common. However, upon silver challenge, high-level silver resistance was selected at high frequency in 76% of isolates of *Enterobacter spp.*, 58% of isolates of *Klebsiella spp.*, and 0.7% of isolates of *E. coli*. Other studies have shown that, although genes encoding silver resistance can be identified in a number of bacterial species, phenotypic resistance is relatively low. Silver resistance was examined in three clinically important *Enterobacteriaceae* genera (Sütterlin et al, 2017). Genes encoding silver resistance were detected most frequently in *Enterobacter spp.* (48%), followed by *Klebsiella spp.* (41%) and *E. coli* 4%. Phenotypical resistance to silver nitrate was found in *Enterobacter* (13%) and *Klebsiella* (3%) isolates. The results of Villapún et al (2021) suggest that the clinical use of silver is unlikely to select for silver resistance. Their studies used three bacteria from the World Health Organization (WHO) priority pathogens list: *E. coli*, *P. aeruginosa* and *S. aureus*. *S. epidermidis* was also studied. And the pathogens were evaluated against silver nitrate.

Several articles report the evolution of silver resistance of several microorganisms when challenged in vitro with AgNPs. McNeilly et al (2023) describe the development of a resistant phenotype in *A. baumannii* to AgNPs. Graves et al (2015), using a non-silver-resistant *E. coli* (using a strain that does not have any known silver resistance elements), found that that exposure to low concentrations of AgNPs resulted in the development of silver resistance, and that this resistance required relatively few mutation steps. A silver-sensitive *P. aeruginosa*, isolated from a burn wound and sensitive to

Figure 3. Review-identified articles published per year and cumulatively related to phenotypic silver resistance in bacteria between 1975 and 2024 ($n=50$).




Figure 3

AgNPs, exhibited reduced silver sensitivity with prolonged exposure to silver; a resistance that was reversed in the subsequent absence of silver (Mann et al, 2021). Stable acquisition of silver resistance has also been identified. Growing *S. aureus* in the presence of sublethal concentrations of AgNPs resulted in the retention of silver resistance in the microorganisms by multiple passages of the bacteria under silver-free conditions (Elbehiry et al, 2019).

The development of silver resistance in microorganisms is heterogeneous; not all microorganisms, when exposed to prolonged periods develop silver resistance. Mutations for silver resistance develop due to selection pressures exerted by silver-based antimicrobials, leading to the emergence of bacteria with mechanisms to tolerate or overcome silver's toxicity. Bacteria mutation rates are high, typically ranging from 10⁶ to 10⁹ base substitutions per nucleotide per generation, but bacteria have been identified with approximately 100-fold higher mutation frequencies in clinical environments (Chevallereau et al, 2019). Natural genetic variation and the selective pressure exerted by prolonged exposure to silver results in heterogeneity in resistance (Habboush and Guzman, 2025).

Is silver resistance a rapidly developing issue in wound pathogens?

Antibiotics have been used widely for the treatment of wound infection. An alternate strategy to the use of antibiotics is the use of antimicrobial metals in the form of metal ions (e.g., silver ions) or nanoparticles. With the growing number of studies identifying the emergence of silver-resistant microorganisms, the implications for wounds and wound infection are particularly worrisome. Silver is considered one of the most potent, especially when prepared as nanoparticles (Roman et al, 2020). Silver has become commonplace in the clinical setting (Hussey et al, 2019). Despite this, examples of acquired resistance are rarely reported in the literature (Binsuwaidan et al, 2024).

Chronic wounds, however, provide a unique environment whereby silver dressings may be present in situ over prolonged periods of time, increasing the potential for bacterial phenotypic adaptation (Binsuwaidan et al, 2024). The use of silver-based treatments has especially increased in burn and wound care (Khansa et al, 2019). As a result of increased silver utilisation, questions concerning antimicrobial stewardship (AMS) and fears of widespread silver resistance emerging in clinical bacteria have been raised (Chopra,

2007). The use of AgNPs has been reported to result in the growth of resistant phenotypes (Gunawan et al, 2017; Panáček et al, 2018), thus calling into question the continuous and long-term utility of these nanomaterials in various formulations, wound dressings, medical devices, or other common household items. It is important to note that the WHO has developed a list of antibiotic resistant, global priority pathogens (WHO, 2024) as leading causes of nosocomial infections, collectively referred to as the ESKAPE pathogens (Rice, 2008; Santajit and Indrawattana, 2016). The ESKAPE organisms represent the model archetype of virulent and adaptive bacterial organisms, as they frequently cause severe and chronic disease and some of these pathogens are instrumental in wound infection.

A number of articles have shown the presence of silver resistance in wound isolate microorganisms which reinforce the concern clinicians have that silver resistance in wound care is becoming a significant issue (Bridges et al, 1979; Pirnay et al, 2003; Loh et al, 2009; Woods et al, 2009; Sütterlin et al, 2012; Finley et al, 2015; Hosny et al, 2019; Percival et al, 2019; Safain et al, 2023; Binsuwaidan et al, 2024). Hosny et al (2019), examining 150 clinical isolates from burns and other wounds found 19 silver-resistant bacterial isolates. In a small study of 14 patients with chronic leg ulcers treated for 3 weeks with silver-based dressings, silver dressings had a limited effect on the primary pathogens (Sütterlin et al, 2012). In vitro evolution of silver resistance in these leg ulcer isolates indicated that it took only three weeks for silver resistance to emerge in isolates with silver resistance genes (sil genes). In a separate study, clinical isolates (*E. coli*, *P. aeruginosa*, *S. aureus*) isolated from patients with chronic diabetic foot wounds were assessed for their sensitivity to silver before and after passaging in the presence of ionic silver. Repeated exposure to ionic silver did not result in planktonic phenotypic silver resistance (Binsuwaidan et al, 2024).

However, there were significant changes in ulcer-derived *Pseudomonas* biofilm formation and sensitivity, with increased levels of biofilm formation being seen in this strain when cultured in the presence of silver. There was also an associated reduction in silver susceptibility.

Finley et al (2015) screened 859 clinical isolates (trauma and burn wounds) noted two isolates (*K. pneumonia* and *E. cloacae*) that were able to grow at high levels of silver, and estimated that only 0.2% had clinically significant phenotypic expression of silver resistance. However, they concluded that

these results indicated silver resistance at a level that was capable of greatly reducing, if not negating, the effectiveness of most commercially available silver dressings, compared to non-resistant counterparts of the same species. The authors suggest silver resistance at a level that could significantly impact wound care and the use of silver-based dressings. Binsuwaidan et al (2024) highlighted that silver resistance in microorganisms is complex, particularly as there are no universally accepted guidelines to define phenotypic silver resistance.

Genetic (genotypic) and molecular biology (mechanism of action, MOA) analysis of silver resistance

Genomic-based silver resistance in bacteria is linked to genes and plasmids that code for silver resistance (Terzioğlu et al, 2022). Decreased susceptibility to silver ions/AgNPs has been linked to silver resistance genes encoding, for example, a silver binding protein, as well as additional genes involved in restricting the presence of silver within the microorganism (Maillard et al, 2021; Terzioğlu et al, 2022). Silver resistance can arise without the presence of silver resistance genes (Randall et al, 2015). Therefore, phenotypic silver resistance whereby microorganisms show signs of silver resistance is likely to provide a better indication of clinical impact than the presence or absence of silver resistance genes themselves. A recent review has proposed that the bacterial mechanism of silver resistance involves a combination of interconnected systems: (1) Inducing extracellular silver aggregation and reduction of Ag⁺ to Ag⁰; (2) Preventing silver from entering cells; (3) Efflux of silver in cells; (4) Self-repair of damage (Li and Xu, 2024).

Our review identified 55 articles related to the study of the genetics of silver resistance and/or the mechanism of action, particularly in the identification of genes and genetic elements in the genetic material of microorganisms that may confer silver resistance. Several microorganisms isolated from chronic diabetic foot wounds were identified as containing silver resistance genes (Percival et al, 2008). Two silver-resistant bacteria were identified: *E. cloacae*, a microorganism the authors point out that is rarely implicated as a primary pathogen in chronic wounds. They also find that no wound pathogens (*S. aureus* and *P. aeruginosa*) were found to contain silver-resistant genes. Studies assessing the prevalence of silver resistance genes animal and human wounds found that the presence of a silver-resistant gene did not afford protection to the organism

in the presence of the silver dressing (Loh et al, 2009; Woods et al, 2009). Finley et al (2015) screened 859 clinical isolates (from trauma and burn wounds) and identified 31 that contained at least one silver resistance gene. However, despite having these genes most of the bacteria displayed little or no increase in resistance to ionic silver. Their findings suggest a low prevalence of these genes (3.6% minimum) occurring in hospital isolates and even fewer (0.2%) with clinically significant phenotypic expression of silver resistance. In a previously discussed study, Safain et al (2023) found that, although 65% (101/155) of sil gene-bearing isolates were resistant for silver nitrate, 17% (59/346) of the sil-negative isolates demonstrated phenotypic silver nitrate resistance (Safain et al, 2023).

What are the implications of silver resistance for wound care?

Finley et al (2015), assaying the effectiveness of silver-based wound dressings against silver resistant strains of microorganisms, indicated that silver resistance was at a level that is capable of greatly reducing, if not negating, the effectiveness of most commercially available silver dressings compared with non-resistant counterparts, and that acute emergence of silver resistance would have extensive consequences on wound therapies. As a consequence of the widespread development of microbial resistance to silver it is imperative to identify potential antimicrobial agents that may be used to combat wound infections that minimises or avoids the development of resistance, are not detrimental to healing (i.e., are not cytotoxic to tissues), do not cause adverse effects (e.g., allergic reactions), but have a wide range of antimicrobial activity. In addition, there should be strict monitoring on the use of silver in medical settings with the establishment of an approved standardised method for silver resistance detection (Hosny et al, 2019).

Our review identified several studies related to the development of therapies as alternatives to silver-based antimicrobials to circumvent the development of silver resistance. One study examined the potential of bismuth-based nanoparticles (BiNPs) (Pant et al, 2022). Another suggests that covalent bonding of silver to cyanographene over comes silver resistance to AgNPs (Panáček et al, 2021). Pant et al (2022) showed that BiNPs exhibited a potent broad-spectrum antimicrobial activity against *P. aeruginosa* and *S. aureus*, and were effective against silver resistant bacteria. BiNPs did not appear to contribute to the development of bismuth-resistant phenotypes

after 30 passages of consecutive exposure to sub-lethal doses of NPs. Cyanographene is an efficient covalent trap for silver ions, and strong covalent immobilisation of silver and has potent antibacterial activity (similar to free silver ions), activity in AgNP-resistant bacterial strains, and very low leaching of silver ions or AgNPs (Panáček et al, 2021). A nanohybrid (AgNPs supported on 1nm-thick silicate “platelets”) showed a strong antibacterial activity on all pathogens assessed, including silver-resistant *E. coli* (Su et al, 2011). The authors suggest that the nanohybrid bypasses the usual ionic silver mechanisms on silver-resistant bacteria leading to effective antibacterial activity in silver-resistant *E. coli*.

Conclusion

This review has identified a substantial number of articles that provide evidence that antimicrobial silver resistance is a growing problem in healthcare. In the general healthcare landscape, a wide variety of pathogens have been identified that have developed silver resistance [Table 5]. The implications of this resistance are serious in that silver is widely used in healthcare, domestic and agricultural applications. Additionally, and importantly, silver treatments are widely used in the prevention/treatment of infection in wounds as dressings, irrigation/instillation solutions and gels etc., to great effect, with an initial two-weeks treatment recommendation followed by a management re-evaluation (Leaper, 2012). But several authors have described wound pathogens as developing resistance, and thus becoming difficult to treat when infection occurs.

Ultimately, the impact of this development of resistance is that these treatments become less effective and the toolbox available to clinicians diminishes. Therefore, while silver remains a valuable tool in wound care, its use should be cautious and accompanied by ongoing research into resistance mechanisms and alternative treatments. The development of new antimicrobial agents that rely upon chemical mechanisms (e.g., bismuth) may also result in the development of resistance. A focus on a physical mechanism-based approach to antimicrobial action may avoid chemical-based resistance mechanisms in the future.

Combining antimicrobial agents with a targeted and sustainable approach could help mitigate the risks associated with resistance and ensure effective wound management. Because of the widespread development of silver resistance, resistance monitoring should form part of AMS practices.

Also, it is imperative that alternative treatments to antibiotics and some antiseptics (e.g., silver) are identified and used as alternatives. These alternatives should have a good in vitro and in vivo/clinical evidence base, not cause any detriment to healing and be cost-effective. In addition, it is important to identify novel antimicrobial agents that do not cause resistance but, at the same time, have wide ranging antimicrobial activity. DACC, which acts via a physical-based mechanism which does not lead to the development of resistance, is an example of an antimicrobial agent used in wound dressings that is today providing a route to delivering effective antimicrobial activity with no resistance. ●

References

- Alhajjar RK, Roche KM, Techtmann SM (2022) Comparative analysis of the mechanism of resistance to silver nanoparticles and the biocide 2,2-dibromo-3-nitripropionamide. *Antimicrob Agents Chemother* 66: e0203121.
- Ali MM, Provoost A, Mijndonckx K, Van Houdt R, Charlier D (2020) DNA-binding and transcription activation by unphosphorylated response regulator AgrR from *Cupriavidus metallidurans* involved in silver resistance. *Front Microbiol* 11: 1635
- Alotaibi AM, Alsaleh NB, Aljasham AT et al (2022) Silver nanoparticle-based combinations with antimicrobial agents against antimicrobial-resistant clinical isolates. *Antibiotics (Basel)* 11(9): 1219
- Andrade LN, Siqueira TES, Martinez R, Darini ALC (2018) Multidrug-resistant CTX-M-(15, 9, 2)- and KPC-2-producing *Enterobacter hormaechei* and *Enterobacter asburiae* isolates possessed a set of acquired heavy metal tolerance genes including a chromosomal sil operon (for acquired silver resistance). *Front Microbiol* 9: 539
- Annear DI, Mee BJ, Bailey M (1976) Instability and linkage of silver resistance, lactose fermentation, and colony structure in *Enterobacter cloacae* from burn wounds. *J Clin Pathol* 29(5): 441-3
- Arrault C, Monneau YR, Martin M et al (2023) The battle for silver binding: How the interplay between the SilE, SilF, and SilB proteins contributes to the silver efflux pump mechanism. *J Biol Chem* 299(8): 105004
- Arrieta-Gisasola A, Martínez-Ballesteros I, Martínez-Malaxetxebarria I et al (2025) Detection of mobile genetic elements conferring resistance to heavy metals in *Salmonella* 4,[5],12:i:- and *Salmonella* Typhimurium serovars and their association with antibiotic resistance. *Int J Food Microbiol* 426: 110890
- Ashraf S, Chatha MA, Ejaz W et al (2014) Lysozyme-coated silver nanoparticles for differentiating bacterial strains on the basis of antibacterial activity. *Nanoscale Res Lett* 9(1): 565
- Asiani KR, Williams H, Bird L et al (2016) SilE is an intrinsically disordered periplasmic “molecular sponge” involved in bacterial silver resistance. *Mol Microbiol* 101(5): 731-42
- Ayobami O, Brinkwirth S, Eckmanns T, Markwart R (2022) Antibiotic resistance in hospital-acquired ESKAPE-E infections in low- and lower-middle-income countries: a systematic review and meta-analysis. *Emerg Microbes Infect* 11(1): 443-51
- Babel L, Nguyen MH, Mittelheisser C et al (2021) NMR reveals the interplay between SilE and SilB model peptides in the context of silver resistance. *Chem Commun (Camb)* 57(70): 8726-9
- Babu MM, Sridhar J, Gunasekaran P (2011) Global

- transcriptome analysis of *Bacillus cereus* ATCC 14579 in response to silver nitrate stress. *J Nanobiotechnology* 9: 49
- Barreto R, Barrois B, Lambert J et al (2020) Addressing the challenges in antiseptics: focus on povidone iodine. *Int J Antimicrob Agents* 56(3): 106064
- Bearson BL, Trachsel JM, Shippy DC et al (2020) The role of *Salmonella* Genomic Island 4 in metal tolerance of *Salmonella enterica* serovar I 4,[5],12:i:- pork outbreak isolate USDA15WA-1. *Genes (Basel)* 11(11): 1291
- Bereanu AS, Bereanu R, Mohor C et al (2024) Prevalence of infections and antimicrobial resistance of ESKAPE group bacteria isolated from patients admitted to the intensive care unit of a county emergency hospital in Romania. *Antibiotics (Basel)* 13(5): 400
- Billman-Jacobe H, Liu Y, Haïtes R et al (2018) pSTM6-275, a conjugative IncHI2 plasmid of *Salmonella enterica* that confers antibiotic and heavy-metal resistance under changing physiological conditions. *Antimicrob Agents Chemother* 62(5): e02357-17
- Binsuwaïdan R, Almuzaini O, Mercer S et al (2024) Variable effects of exposure to ionic silver in wound-associated bacterial pathogens. *Lett Appl Microbiol* 77(4): ova030
- Blackburn J, Ousey K, Patton D et al (2023) What is the evidence that there is antimicrobial resistance associated with the use of topical antimicrobial preparations? *Wound Practice Res* 31(1): 40-8
- Blanco Massani M, Klumpp J, Widmer M et al (2018) Chromosomal Sil system contributes to silver resistance in *E. coli* ATCC 8739. *Biometals* 31(6): 1101-14
- Brady MJ, Lisay CM, Yurkovetskiy AV, Sawan SP (2003) Persistent silver disinfectant for the environmental control of pathogenic bacteria. *Am J Infect Control* 31(4): 208-14
- Bridges K, Kidson A, Lowbury EJ, Wilkins MD (1979) Gentamicin- and silver-resistant *Pseudomonas* in a burns unit. *Br Med J* 1(6161): 446-9
- Chabert V, Hologne M, Sèneque O et al (2018) Alpha-helical folding of SilE models upon Ag(His)(Met) motif formation. *Chem Commun (Camb)* 54(74): 10419-22
- Chalmers G, Rozas KM, Amachawadi RG et al (2018) Distribution of the *pco* gene cluster and associated genetic determinants among swine *Escherichia coli* from a controlled feeding trial. *Genes (Basel)* 9(10): 504
- Chevallereau A, Meaden S, van Houte S et al (2019) The effect of bacterial mutation rate on the evolution of CRISPR-Cas adaptive immunity. *Philos Trans R Soc Lond B Biol Sci* 374(1772): 20180094
- Chopra I (2007) The increasing use of silver-based products as antimicrobial agents: a useful development or a cause for concern? *J Antimicrob Chemother* 59(4): 587-90
- Cidre I, Pulido RP, Burgos MJG et al (2017) Copper and zinc tolerance in bacteria isolated from fresh produce. *J Food Prot* 80(6): 969-75
- Clark CG, Landgraaf C, Robertson J et al (2020) Distribution of heavy metal resistance elements in Canadian *Salmonella* 4,[5],12:i:- populations and association with the monophasic genotypes and phenotype. *PLoS One* 15(7): e0236436
- Corona F, Martinez J (2013) Phenotypic resistance to antibiotics. *Antibiotics (Basel)* 2(2): 237-55
- Doody O, Bailey ME (2016) Setting a research question, aim and objective. *Nurse Res* 23(4): 19-23
- Dunne KA, Chaudhuri RR, Rossiter AE et al (2017) Sequencing a piece of history: complete genome sequence of the original *Escherichia coli* strain. *Microb Genom* 3(3): mgen000106
- Elarabi NI, Halema AA, Abdelhadi AA et al (2023) Draft genome of *Raoultella planticola*, a high lead resistance bacterium from industrial wastewater. *AMB Express* 13(1): 14
- Elbehiry A, Al-Dubaib M, Marzouk E, Moussa I (2019) Antibacterial effects and resistance induction of silver and gold nanoparticles against *Staphylococcus aureus*-induced mastitis and the potential toxicity in rats. *Microbiologyopen* 8(4): e00698
- Elkrewi E, Randall CP, Ooi N et al (2017) Cryptic silver resistance is prevalent and readily activated in certain Gram-negative pathogens. *J Antimicrob Chemother* 72(11): 3043-6
- Ferdous Z, Nemmar A (2020) Health impact of silver nanoparticles: a review of the biodistribution and toxicity following various routes of exposure. *Int J Mol Sci* 21(7): 2375
- Finley PJ, Norton R, Austin C et al (2015) Unprecedented silver resistance in clinically isolated Enterobacteriaceae: major implications for burn and wound management. *Antimicrob Agents Chemother* 59(8): 4734-41
- Foka FET, Mienie C, Bezuidenhout CC, Ateba CN (2020) Complete genomic analysis of VRE from a cattle feedlot: focus on 2 antibiotic resistance. *Front Microbiol* 11: 571958.
- Franke S, Grass G, Nies DH (2001) The product of the *ybdE* gene of the *Escherichia coli* chromosome is involved in detoxification of silver ions. *Microbiology (Reading)* 147 (Pt 4): 965-72
- Frei A, Verderosa AD, Elliott AG et al (2023) Metals to combat antimicrobial resistance. *Nat Rev Chem* 7(3):202-24.
- Fuentes-Castillo D, Navas-Suárez PE, Gondim MF et al (2021) Genomic characterization of multidrug-resistant ESBL-producing *Escherichia coli* ST58 causing fatal colibacillosis in critically endangered Brazilian merganser (*Mergus octosetaceus*). *Transbound Emerg Dis* 68(2): 258-66
- Gillieatt BF, Coleman NV (2024) Unravelling the mechanisms of antibiotic and heavy metal resistance co-selection in environmental bacteria. *FEMS Microbiol Rev* 48(4): fue017
- Gokulan K, Williams K, Khare S (2017) Silver ion-mediated killing of a food pathogen: melting curve analysis data of silver resistance genes and growth curve data. *Data Brief* 11: 49-53
- González-Fernández S, Lozano-Iturbe V, García B et al (2020) Antibacterial effect of silver nanorings. *BMC Microbiol* 20(1): 172
- Graves JL Jr, Tajkarimi M, Cunningham Q et al (2015) Rapid evolution of silver nanoparticle resistance in *Escherichia coli*. *Front Genet* 6: 42
- Grim CJ, Kotewicz ML, Power KA et al (2013) Pan-genome analysis of the emerging foodborne pathogen *Cronobacter* spp. suggests a species-level bidirectional divergence driven by niche adaptation. *BMC Genomics* 14: 366
- Gugala N, Lemire J, Chatfield-Reed K et al (2018) Using a chemical genetic screen to enhance our understanding of the antibacterial properties of silver. *Genes (Basel)* 9(7): 344
- Gugala N, Vu D, Parkins MD, Turner RJ (2019) Specificity in the susceptibilities of *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* clinical isolates to six metal antimicrobials. *Antibiotics (Basel)* 8(2): 51
- Gullberg E, Albrecht LM, Karlsson C et al (2014) Selection of a multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. *mBio* 5(5): e01918-14
- Gunawan C, Marquis CP, Amal R et al (2017) Widespread and indiscriminate nanosilver use: genuine potential for microbial resistance. *ACS Nano* 11(4): 3438-45
- Gunawan C, Teoh WY, Marquis CP, Amal R (2013) Induced adaptation of *Bacillus* sp. to antimicrobial nanosilver. *Small* 9(21): 3554-60
- Guo J, Qin S, Wei Y et al (2019) Silver nanoparticles exert concentration-dependent influences on biofilm development and architecture. *Cell Prolif* 52(4): e12616
- Gupta A, Matsui K, Lo JF, Silver S (1999) Molecular basis for resistance to silver cations in *Salmonella*. *Nat Med* 5(2): 183-8
- Gupta A, Maynes M, Silver S (1998) Effects of halides on plasmid-mediated silver resistance in *Escherichia coli*. *Appl Environ Microbiol* 64(12): 5042-5
- Gupta A, Phung LT, Taylor DE, Silver S (2001) Diversity of silver resistance genes in IncH incompatibility group plasmids. *Microbiology (Reading)* 147 (pt 12): 3393-402
- Gupta A (1999) RT-PCR: characterization of long multi-gene operons and multiple transcript gene clusters in bacteria. *Biotechniques* 27(5): 966-70, 972

- Habboush Y, Guzman N (2025) *Antibiotic Resistance*. [Updated 2023 Jun 20]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK513277/> (accessed 17.04.2025)
- Haefeli C, Franklin C, Hardy K (1984) Plasmid-determined silver resistance in *Pseudomonas stutzeri* isolated from a silver mine. *J Bacteriol* 158(1): 389–92
- Håkonsholm F, Hetland MAK, Löhr IH et al (2023) Co-localization of clinically relevant antibiotic- and heavy metal resistance genes on plasmids in *Klebsiella pneumoniae* from marine bivalves. *Microbiologypopen* 12(4): e1368
- Håkonsholm F, Hetland MAK, Svanevik CS et al (2022) Insights into the genetic diversity, antibiotic resistance and pathogenic potential of *Klebsiella pneumoniae* from the Norwegian marine environment using whole-genome analysis. *Int J Hyg Environ Health* 242: 113967
- Hanczvikkel A, Füzi M, Ungvári E, Tóth Á (2018) Transmissible silver resistance readily evolves in high-risk clone isolates of *Klebsiella pneumoniae*. *Acta Microbiol Immunol Hung* 65(3): 387–403
- He J, Du X, Zeng X et al (2022) Phenotypic and genotypic characterization of a hypervirulent carbapenem-resistant *Klebsiella pneumoniae* ST17–KL38 clinical isolate harboring the carbapenemase IMP-4. *Microbiol Spectr* 10(2): e0213421
- Hosnedlova B, Kabanov D, Kepinska M et al (2022) Effect of biosynthesized silver nanoparticles on bacterial biofilm changes in *S. aureus* and *E. coli*. *Nanomaterials (Basel)* 12(13): 2183
- Hosny AEM, Rasmy SA, Aboul-Magd DS et al (2019) The increasing threat of silver-resistance in clinical isolates from wounds and burns. *Infect Drug Resist* 12(13): 1985–2001
- Hussey L, Stocks SJ, Wilson P et al (2019) Use of antimicrobial dressings in England and the association with published clinical guidance: interrupted time series analysis. *BMJ Open* 9(9): e028727
- Jangid H, Singh S, Kashyap P et al (2024) Advancing biomedical applications: an in-depth analysis of silver nanoparticles in antimicrobial, anticancer, and wound healing roles. *Front Pharmacol* 15: 1438227
- Johnson TJ, Siek KE, Johnson SJ, Nolan LK (2005) DNA sequence and comparative genomics of pAPEC-O2-R, an avian pathogenic *Escherichia coli* transmissible R plasmid. *Antimicrob Agents Chemother* 49(11): 4681–8
- Joseph S, Desai P, Ji Y et al (2012) Comparative analysis of genome sequences covering the seven *Cronobacter* species. *PLoS One* 7(11): e49455
- Kale SK, Parishwad GV, Patil AS (2021) Emerging agriculture applications of silver nanoparticles. *ES Food Agrofor* 3: 17–22
- Kamat S, Kumari M (2023) Emergence of microbial resistance against nanoparticles: mechanisms and strategies. *Front Microbiol* 14: 1102615
- Kamatheewatta K, Bushell R, Rafa F et al (2020) Colonization of a hand washing sink in a veterinary hospital by an *Enterobacter hormaechei* strain carrying multiple resistances to high importance antimicrobials. *Antimicrob Resist Infect Control* 9(1): 163
- Kaur P, Vadehra DV (1986) Mechanism of resistance to silver ions in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 29(1): 165–7
- Kędziora A, Wernecki M, Korzekwa K et al (2020) Consequences of long-term bacteria's exposure to silver nanoformulations with different physicochemical properties. *Int J Nanomedicine* 15: 199–213
- Khansa I, Schoenbrunner AR, Kraft CT, Janis JE (2019) Silver in wound care—friend or foe?: A comprehensive review. *Plast Reconstr Surg Glob Open* 7(8): e2390
- Khor SY, Jegathesan M (1983) Heavy metal and disinfectant resistance in clinical isolates of gram-negative rods. *Southeast Asian J Trop Med Public Health* 14(2): 199–203
- Klonowska A, Moulin L, Ardley JK et al (2020) Novel heavy metal resistance gene clusters are present in the genome of *Cupriavidus neocaledonicus* STM 6070, a new species of *Mimosa pudica* microsymbiont isolated from heavy-metal-rich mining site soil. *BMC Genomics* 21(1): 214
- Kremer AN, Hoffmann H (2012) Subtractive hybridization yields a silver resistance determinant unique to nosocomial pathogens in the *Enterobacter cloacae* complex. *J Clin Microbiol* 50(10): 3249–57
- Kucerova E, Clifton SW, Xia XQ et al (2010) Genome sequence of *Cronobacter sakazakii* BAA-894 and comparative genomic hybridization analysis with other *Cronobacter* species. *PLoS One* 5(3): e9556
- Lansdown AB (2006) Silver in health care: antimicrobial effects and safety in use. *Curr Probl Dermatol* 33: 17–34
- Lau CP, Abdul-Wahab MF, Jaafar J et al (2017) Toxic effect of high concentration of sonochemically synthesized polyvinylpyrrolidone-coated silver nanoparticles on *Citrobacter* sp. A1 and *Enterococcus* sp. C1. *J Microbiol Immunol Infect* 50(4): 427–34
- Leaper D (2012) Appropriate use of silver dressings in wounds: international consensus document. *Int Wound J* 9(5): 461–4
- Li H, Xu H (2024) Mechanisms of bacterial resistance to environmental silver and antimicrobial strategies for silver: a review. *Environ Res* 248: 118313
- Li XZ, Nikaido H, Williams KE (1997) Silver-resistant mutants of *Escherichia coli* display active efflux of Ag⁺ and are deficient in porins. *J Bacteriol* 179(19): 6127–32
- Liao S, Zhang Y, Pan X et al (2019) Antibacterial activity and mechanism of silver nanoparticles against multidrug-resistant *Pseudomonas aeruginosa*. *Int J Nanomedicine* 14: 1469–87
- Lima de Silva AA, de Carvalho MA, de Souza SA et al (2012) Heavy metal tolerance (Cr, Ag and Hg) in bacteria isolated from sewage. *Braz J Microbiol* 43(4): 1620–31
- Loh JV, Percival SL, Woods EJ et al (2009) Silver resistance in MRSA isolated from wound and nasal sources in humans and animals. *Int Wound J* 6: 32–8
- Losasso C, Belluco S, Cibin V et al (2014) Antibacterial activity of silver nanoparticles: sensitivity of different *Salmonella* serovars. *Front Microbiol* 5: 227
- Maillard JY, Kampf G, Cooper R (2021) Antimicrobial stewardship of antiseptics that are pertinent to wounds: the need for a united approach. *JAC Antimicrob Resist* 3(1): dlab027
- Mann R, Holmes A, McNeilly O et al (2021) Evolution of biofilm-forming pathogenic bacteria in the presence of nanoparticles and antibiotic: adaptation phenomena and cross-resistance. *J Nanobiotechnology* 19(1): 291
- Mastrolilli E, Pietrucci D, Barco L et al (2018) A comparative genomic analysis provides novel insights into the ecological success of the monophasic *Salmonella* serovar 4,[5],12:i: *Front Microbiol* 9: 715
- May A, Kopecki Z, Carney B, Cowin A (2022) Antimicrobial silver dressings: a review of emerging issues for modern wound care. *ANZ J Surg* 92(3): 379–84
- McCarlie SJ, Boucher CE, Bragg RR (2023) Genomic islands identified in highly resistant *Serratia* sp. HRI: a pathway to discover new disinfectant resistance elements. *Microorganisms* 11(2): 515
- McHugh GL, Moellering RC, Hopkins CC, Swartz MN (1975) *Salmonella typhimurium* resistant to silver nitrate, chloramphenicol, and ampicillin. *Lancet* 1(7901): 235–40
- McNeilly O, Mann R, Cummins ML et al (2023) Development of nanoparticle adaptation phenomena in *Acinetobacter baumannii*: physiological change and defense response. *Microbiol Spectr* 11(1): e0285722
- McNeilly O, Mann R, Hamidian M, Gunawan C (2021) Emerging concern for silver nanoparticle resistance in *Acinetobacter baumannii* and other bacteria. *Front Microbiol* 12: 652863
- Miloud SB, Dziri O, Ferjani S et al (2021) First description of various bacteria resistant to heavy metals and antibiotics isolated from polluted sites in Tunisia. *Pol J Microbiol* 70(2): 161–74
- Monych NK, Turner RJ (2020) Multiple compounds secreted by *Pseudomonas aeruginosa* increase the tolerance

- of *Staphylococcus aureus* to the antimicrobial metals copper and silver. *mSystems* 5(5): e00746-20
- Muller M, Merrett ND (2014) Pyocyanin production by *Pseudomonas aeruginosa* confers resistance to ionic silver. *Antimicrob Agents Chemother* 58(9): 5492-9
- Muller M (2018) Bacterial silver resistance gained by cooperative interspecies redox behavior. *Antimicrob Agents Chemother* 62(8): e00672-18
- Nicolás MF, Ramos PIP, Marques de Carvalho F et al (2018) Comparative genomic analysis of a clinical isolate of *Klebsiella quasipneumoniae* subsp. *similipneumoniae*, a KPC-2 and OKP-B-6 beta-lactamases producer harboring two drug-resistance plasmids from Southeast Brazil. *Front Microbiol* 9: 220
- Norton R, Finley PJ (2021) Clinically isolated bacteria resistance to silver-based wound dressings. *J Wound Care* 30(3): 238-47
- Panáček A, Kvítek L, Smékalová M et al (2018) Bacterial resistance to silver nanoparticles and how to overcome it. *Nat Nanotechnol* 13(1): 65-71
- Panáček D, Hochvaldová L, Bakandritsos A et al (2021) Silver covalently bound to cyanographene overcomes bacterial resistance to silver nanoparticles and antibiotics. *Adv Sci (Weinh)* 8(12): 2003090
- Pant BD, Benin BM, Abeydeera N et al (2022) Bi2O3 nanoparticles exhibit potent broad-spectrum antimicrobial activity and the ability to overcome Ag-, ciprofloxacin- and meropenem-resistance in *P. aeruginosa*: the next silver bullet of metal antimicrobials? *Biomater Sci* 10(6): 1523-31
- Pelgrift RY, Friedman AJ (2013) Nanotechnology as a therapeutic tool to combat microbial resistance. *Adv Drug Deliv Rev* 65(13-14): 1803-15.
- Percival SL, Salisbury AM, Chen R (2019) Silver, biofilms and wounds: resistance revisited. *Crit Rev Microbiol* 45(2): 223-37
- Percival SL, Thomas J, Linton S et al (2012) The antimicrobial efficacy of silver on antibiotic-resistant bacteria isolated from burn wounds. *Int Wound J* 9(5): 488-93
- Percival SL, Woods E, Nutekpor M et al (2008) Prevalence of silver resistance in bacteria isolated from diabetic foot ulcers and efficacy of silver-containing wound dressings. *Ostomy Wound Manage* 54(3): 30-40
- Pirnay JP, De Vos D, Cochez C et al (2003) Molecular epidemiology of *Pseudomonas aeruginosa* colonization in a burn unit: persistence of a multidrug-resistant clone and a silver sulfadiazine-resistant clone. *J Clin Microbiol* 41(3): 1192-202
- Potgieter MD, Meidany P (2018) Evaluation of the penetration of nanocrystalline silver through various wound dressing mediums: an in vitro study. *Burns* 44(3): 596-602
- Randall CP, Gupta A, Jackson N et al (2015) Silver resistance in Gram-negative bacteria: a dissection of endogenous and exogenous mechanisms. *J Antimicrob Chemother* 70(4): 1037-46
- Rice LB (2008) Federal funding for the study of antimicrobial resistance in nosocomial pathogens: no ESKAPE. *J Infect Dis* 197(8): 1079-81
- Riley TV, Mee BJ (1982) Susceptibility of *Bacteroides* spp. to heavy metals. *Antimicrob Agents Chemother* 22(5): 889-92
- Rippon M, Rogers AA, Ousey K, Chadwick P (2023) Experimental and clinical evidence for DACC-coated dressings: an update. *J Wound Care* 32(Sup8a): S13-22
- Rippon MG, Rogers AA, Ousey K (2021) Antimicrobial stewardship strategies in wound care: evidence to support the use of dialkylcarbamoyl chloride (DACC)-coated wound dressings. *J Wound Care* 30(4): 284-96
- Rippon MG, Rogers AA, Ousey K (2023) Polyhexamethylene biguanide and its antimicrobial role in wound healing: a narrative review. *J Wound Care* 32(1): 5-20
- Rodrigues AS, Batista JGS, Rodrigues MÁV et al (2024) Advances in silver nanoparticles: a comprehensive review on their potential as antimicrobial agents and their mechanisms of action elucidated by proteomics. *Front Microbiol* 15: 1440065
- Roman M, Rigo C, Castillo-Michel H et al (2020) Spatiotemporal distribution and speciation of silver nanoparticles in the healing wound. *Analyst* 145(20): 6456-69
- Rybka M, Mazurek Ł, Konop M (2022) Beneficial effect of wound dressings containing silver and silver nanoparticles in wound healing—from experimental studies to clinical practice. *Life (Basel)* 13(1): 69
- Saeb ATM, Al-Rubeaan KA, Abouelhoda M et al (2017) Genome sequencing and analysis of the first spontaneous Nanosilver resistant bacterium *Proteus mirabilis* strain SCDRI. *Antimicrob Resist Infect Control* 6: 119
- Safain KS, Islam MS, Amatullah J et al (2023) Prevalence of silver resistance determinants and extended-spectrum β -lactamases in bacterial species causing wound infection: First report from Bangladesh. *New Microbes New Infect* 52: 101104
- Salomoni R, Léo P, Montemor AF et al (2017) Antibacterial effect of silver nanoparticles in *Pseudomonas aeruginosa*. *Nanotechnol Sci Appl* 10: 115-21
- Sanjar F, Millan CP, Leung KP (2024) Phylogenetic evaluation and genotypic identification of burn-related *Pseudomonas aeruginosa* strains isolated from post-burn human infections during hospitalization. *Pathog Dis* 82: ftac021
- Sano E, Fontana H, Esposito F et al (2023) Genomic analysis of fluoroquinolone-resistant *Leclercia adecarboxylata* carrying the ISKpn19-orf-qnrS1- Δ IS3-blaLAP-2 module in a synanthropic pigeon, Brazil. *J Glob Antimicrob Resist* 33: 256-9
- Santajit S, Indrawattana N (2016) Mechanisms of antimicrobial resistance in ESKAPE pathogens. *Biomed Res Int* 2016: 2475067
- Scepankova H, Combarros-Fuertes P, Fresno JM et al (2021) Role of honey in advanced wound care. *Molecules* 26(16): 4784
- Schultz G, Bjarnsholt T, James GA et al (2017) Consensus guidelines for the identification and treatment of biofilms in chronic nonhealing wounds. *Wound Repair Regen* 25(5): 744-57
- Sedlak RH, Hnilova M, Grosh C et al (2012) Engineered *Escherichia coli* silver-binding periplasmic protein that promotes silver tolerance. *Appl Environ Microbiol* 78(7): 2289-96
- Silver S, Gupta A, Matsui K, Lo JF (1999) Resistance to Ag(I) cations in bacteria: environments, genes and proteins. *Met Based Drugs* 6(4-5): 315-20
- Souza SSR, Turcotte MR, Li J et al (2022) Population analysis of heavy metal and biocide resistance genes in *Salmonella enterica* from human clinical cases in New Hampshire, United States. *Front Microbiol* 13: 983083
- Stabryla LM, Johnston KA, Diemler NA et al (2021) Role of bacterial motility in differential resistance mechanisms of silver nanoparticles and silver ions. *Nat Nanotechnol* 16(9): 996-1003
- Staehlin BM, Gibbons JG, Rokas A et al (2016) Evolution of a heavy metal homeostasis/resistance island reflects increasing copper stress in Enterobacteria. *Genome Biol Evol* 8(3): 811-26
- Su HL, Lin SH, Wei JC et al (2011) Novel nanohybrids of silver particles on clay platelets for inhibiting silver-resistant bacteria. *PLoS One* 6(6): e21125
- Summer M, Ali S, Tahir HM et al (2024) Mode of action of biogenic silver, zinc, copper, titanium and cobalt nanoparticles against antibiotics resistant pathogens. *J Inorg Organomet Polymers Mater* 34(4): 1417-51
- Sütterlin S, Dahlöf M, Tellgren-Roth C et al (2017) High frequency of silver resistance genes in invasive isolates of *Enterobacter* and *Klebsiella* species. *J Hosp Infect* 96(3): 256-61
- Sütterlin S, Edquist P, Sandegren L et al (2014) Silver resistance genes are overrepresented among *Escherichia coli* isolates with CTX-M production. *Appl Environ Microbiol* 80(22): 6863-9

- Sütterlin S, Tano E, Bergsten A et al (2012) Effects of silver-based wound dressings on the bacterial flora in chronic leg ulcers and its susceptibility in vitro to silver. *Acta Derm Venereol* 92(1): 34–9
- Sütterlin S, Téllez-Castillo CJ, Anselem L et al (2018) Heavy metal susceptibility of *Escherichia coli* isolated from urine samples from Sweden, Germany, and Spain. *Antimicrob Agents Chemother* 62(5): e00209–18
- Terzioğlu E, Arslan M, Balaban BG, Çakar ZP (2022) Microbial silver resistance mechanisms: recent developments. *World J Microbiol Biotechnol* 38(9): 158
- Totty JP, Bua N, Smith GE et al (2017) Dialkylcarbamoyl chloride (DACC)-coated dressings in the management and prevention of wound infection: a systematic review. *J Wound Care* 26(3): 107–14
- Valentin E, Bottomley AL, Chilambi GS et al (2020) Heritable nanosilver resistance in priority pathogen: a unique genetic adaptation and comparison with ionic silver and antibiotics. *Nanoscale* 12(4): 2384–92
- Vasishtha R, Chhibber S, Saxena M (1989) Heavy metal resistance in clinical isolates of *Pseudomonas aeruginosa*. *Folia Microbiol (Praha)* 34(5): 448–52
- Vázquez X, García-Fierro R, Fernández J et al (2023) Incidence and genomic background of antibiotic resistance in food-borne and clinical isolates of *Salmonella enterica* serovar Derby from Spain. *Antibiotics* (Basel) 12(7): 1204
- Vilela FP, Rodrigues DDP, Allard MW, Falcão JP (2022) Prevalence of efflux pump and heavy metal tolerance encoding genes among *Salmonella enterica* serovar Infantis strains from diverse sources in Brazil. *PLoS One* 17(11): e0277979
- Villapún VM, Balacco DL, Webber MA et al (2021) Repeated exposure of nosocomial pathogens to silver does not select for silver resistance but does impact ciprofloxacin susceptibility. *Acta Biomater* 134: 760–73
- Wang H, Li J, Min C et al (2022) Characterization of silver resistance and coexistence of sil operon with antibiotic resistance genes among Gram-negative pathogens isolated from wound samples by using whole-genome sequencing. *Infect Drug Resist* 15: 1425–37
- Wiegand C, Abel M, Ruth P, Hipler UC (2012) Analysis of the adaptation capacity of *Staphylococcus aureus* to commonly used antiseptics by microplate laser nephelometry. *Skin Pharmacol Physiol* 25(6): 288–97
- Woods EJ, Cochrane CA, Percival S (2009) Prevalence of silver resistance genes in bacteria isolated from human and horse wounds. *Vet Microbiol* 138(3–4): 325–9
- Woolley CA, Sutton JM, Wand ME (2022) Mutations in SilS and CusS/OmpC represent different routes to achieve high level silver ion tolerance in *Klebsiella pneumoniae*. *BMC Microbiol* 22(1): 113
- World Health Organization (2024) *WHO Bacterial Priority Pathogens List*. Available at: <https://www.who.int/publications/i/item/9789240093461> (accessed 13.03.2025)
- Wound Infection in Clinical Practice Expert Working Group (2008) Wound infection in clinical practice. An international consensus. *Int Wound J* 5 Suppl 3: iii–11
- Woyda R, Oladeinde A, Endale D et al (2024) Genetic characteristics of *Salmonella* isolates recovered from reused broiler litter over three successive flocks. *J Food Prot* 87(3): 100236
- Wu K, Li H, Cui X et al (2022) Mutagenesis and resistance development of bacteria challenged by silver nanoparticles. *Antimicrob Agents Chemother* 66(10): e0062822
- Wu MY, Suryanarayanan K, van Ooij WJ, Oerther DB (2007) Using microbial genomics to evaluate the effectiveness of silver to prevent biofilm formation. *Water Sci Technol* 55(8–9): 413–9
- Yang H, Wei SH, Hobman JL, Dodd CER (2020) Antibiotic and metal resistance in *Escherichia coli* isolated from pig slaughterhouses in the United Kingdom. *Antibiotics* (Basel) 9(11): 746
- Yin IX, Zhang J, Zhao IS et al (2020) The antibacterial mechanism of silver nanoparticles and its application in dentistry. *Int J Nanomedicine* 15: 2555–62
- Yousefian F, Hesari R, Jensen T et al (2023) Antimicrobial wound dressings: a concise review for clinicians. *Antibiotics* (Basel) 12(9): 1434
- Zhao X, Kuipers OP (2021) Synthesis of silver-nisin nanoparticles with low cytotoxicity as antimicrobials against biofilm-forming pathogens. *Colloids Surf B Biointerfaces* 206: 111965
- Zingale GA, Oliveri V, Grasso G (2023) Insights into the binding of Ag ions with SilE model peptides: an NMR and MS coupled approach. *Metallomics* 15(4): mfad015
- Zingali T, Chapman TA, Webster J et al (2020) Genomic characterisation of a multiple drug resistant IncHI2 ST4 plasmid in *Escherichia coli* ST744 in Australia. *Microorganisms* 8(6): 896