

Leukoplast STEPS

The complete care solution for skin tears **#SaveOurSkin**

PREVENT

Skin tear incidence
can be reduced by 50%
when a moisturizer is applied
twice-daily.¹

TREATMENT

What do skin tears look like?



Type 1
No skin loss

Type 2
Partial flap loss

Type 3
Total flap loss

Steps of treatment²

- Control bleeding
- Cleanse and debride
- Manage infection/inflammation
- Consider moisture balance/exudate control
- Monitor wound edge/closure
- Ensure skin flap remains undisturbed
- Depending on the clinical conditions of the wound, the dressing may remain in place for up to 7 days³

RECOVERY

Steps of treatment²

KEEP IT CLEAN:
TENA ProSkin® Dry Washcloths are a soft, easy to use alternative to reusable washcloths. For effective, yet gentle cleansing of fecal residues, use with TENA ProSkin® Cleansing Cream.

KEEP IT PROTECTED:
TENA ProSkin® Protective Cream protects skin from irritation caused by contact with urine and/or stool; helps prevent and relieve chafed, chapped, or cracked skin. TENA ProSkin® Barrier Cream helps treat and prevent skin rashes due to incontinence; soothes chafed skin and helps seal out wetness.


KEEP IT CLEAN:



KEEP IT PROTECTED:





Build your individual solution:



Cuticell® Contact®
low adherent, silicone wound contact layer,
dressing permeable to air and exudate

+



Cutimed® Siltec® Light **Hypafix® skin sensitive**

Or a ready-to-use product:

●●●●-●●●● For low to moderate exudate



Leukoplast®

Leukomed® skin sensitive

●●●●-●●●● For moderate to high exudate



Cutimed®

Cutimed® Siltec® B

KEEP IT CLEAN:



KEEP IT PROTECTED:




COMPRESSION THERAPY:
If the limb shows signs of edema consider applying compression. A full leg assessment should be completed before applying compression.²



Learn more about Leukoplast® STEPS

Skin Tear Effective Protocol and Solutions: Discover STEPS, our complete solution for identifying, preventing, and treating skin tears. We're here to support you as you support your patients.



For Skin Tears on lower limb, please consult healthcare professional for assessment for compression needs.

¹ Carville K, et al. The effectiveness of a twice-daily skin-moisturising regimen for reducing the incidence of skin tears. *International Wound Journal*. 2014;11: 446-53.

² LeBlanc K, et al. Best practice recommendations for the prevention and management of skin tears in aged skin. *Wounds International*. 2018.

³ Deprez J, et al. Measures that patients living in the community can take for the prevention and treatment of skin tears: A comprehensive review of the literature. *J Wound Management*. 2022;23(2):70-77.